

A New Record of *Oniticellus cinctus* (Coleoptera, Scarabaeidae) from Borneo

Masahiro KON¹⁾, Teruo OCHI²⁾, Masahiro ÔHARA³⁾, Akira UEDA⁴⁾ and Sri HARTINI⁵⁾

¹⁾ School of Environmental Science, The University of Shiga Prefecture,
Hassaka-chô 2500, Hikone, 522-8533 Japan

²⁾ Kôfudai 5-21-6, Toyono-chô, Toyono-gun, Osaka, 563-0104 Japan

³⁾ The Hokkaido University Museum, Hokkaido University, Sapporo, 060-0810 Japan

⁴⁾ Hokkaido Research Center, Forestry and Forest Products Research Institute,
Toyohira, Sapporo, 062-8516 Japan

⁵⁾ Zoology Division, Research Center for Biology, LIPI, Bogor, Indonesia

We have been conducting researches on the dung beetle assemblages in East Kalimantan, Indonesia. In the course of our researches, we collected *Oniticellus cinctus* (FABRICIUS). This is the first record of this species from Borneo.

We herewith report the collection data. We referred to BALTHASAR (1963) for the known distribution of this species.

Oniticellus cinctus (FABRICIUS)

Scarabaeus cinctus FABRICIUS, 1775, Syst. Ent., 30.

Specimens examined. 1 ♂, 1 ♀, Sungai Wain, Balikpapan, Kalimantan Timur, Indonesia, 27-XII-2006, A. UEDA leg.; 1 ex., Karya Merdeka RT09, Samboja, Kutai Kartanegara, Kalimantan Timur, Indonesia, 18-XI-2007, M. ÔHARA leg.; 1 ex., Kanpon Timur, Balikpapan, Kalimantan Timur, Indonesia, 18-XI-2007, M. ÔHARA leg.

Distribution. South China, North India, Myanmar, Vietnam, Laos, Thailand, Malay Peninsula, Java, Borneo (new record).

This study was supported in part by a Grant-in-Aid from the Japan Society for the Promotion of Science (No. 17405011) and a grant from the Ministry of the Environment of Japan (Environmental Research by National Research Institute of Government Ministries & Agencies 2004, Evaluation and Forecasting of the CDM-Plantation influences on Biodiversity).

Reference

- BALTHASAR, V., 1963. Monographie der Scarabaeidae und Aphodiidae der palaearktischen und orientalischen Region, 2. 1-628. Tschechoslowakische Akademie der Wissenschaften, Prag.